

AD-A166 055

SIEVES FOR GAUSSIAN PROCESSES(U) WISCONSIN
UNIV-MILWAUKEE DEPT OF MATHEMATICAL SCIENCES J H BEDER
27 NOV 85 AFOSR-TR-86-0069 AFOSR-84-0329


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART

AFOSR-TR- 86-0069

Sieves for Gaussian Processes

DTIC
ELECTE
APR 1 1986

S

D

B

2,3. Research objectives, status of work:

Many phenomena can be modelled as signals to which noise has been added. Frequently, it is assumed that the noise is governed by the so-called Gaussian probability law. If the signal itself is non-random, the most important questions are extraction of the signal and description of the noise. Statistically these are the problems of estimating the mean and covariance, respectively, of a Gaussian process. A classical statistical method, maximum likelihood estimation, breaks down when the signal is "infinite-dimensional," which is the case in many applications. The method of sieves often provides reasonable estimators in such circumstances, and the PI has introduced sieve estimators for the mean and covariance of a Gaussian process. This project is aimed in part at investigating the properties of these estimators, with no assumptions on the "time" parameter (it is only assumed to be an element of a set T).

A sieve estimator is doubly indexed. The first index, say n , is the sample size - in this case, the number of realizations of the process that one observes. The second, say d , is a sieve parameter. Both these indices are allowed to increase without bound to obtain certain asymptotic properties of the estimator; one typically requires that d go to infinity at some rate with respect to n . For a more complete account of sieve estimation, see Grenander [1981] or Geman and Hwang [1982].

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12.
Distribution unlimited.
MATTHEW J. KERRY
Chief, Technical Information Division

Approved for public release;
distribution unlimited.

AD-A166 055
DTIC FILE COPY

The first goal of this project is to develop the properties of a new sieve estimator for the mean, say $m(t)$, of a Gaussian process when the covariance is known. Let \mathcal{H} be the reproducing kernel Hilbert space of the covariance function; then \mathcal{H} is the set of all possible mean functions of the process. Consider the following condition:

(S) the true mean function belongs to a known separable subspace \mathcal{H}_0 of \mathcal{H} .

(If \mathcal{H} is separable, we take $\mathcal{H}_0 = \mathcal{H}$.) The sieve estimator, say $\hat{m}(t)$, is defined in terms of a complete orthonormal basis of \mathcal{H}_0 ; a precise definition of \hat{m} is given in the project proposal. The PI has now shown the following:

a. For fixed n and d , the estimator $\hat{m}(t)$ is a Gaussian process. When (S) holds, one can give a simple formula for the mean and covariance of $\hat{m}(t)$, from which one can show that, at each t , $\hat{m}(t)$ is asymptotically unbiased as $d \rightarrow \infty$ and is weakly consistent for $m(t)$ if $d = O(n)$.

b. Both \hat{m} and m are elements of \mathcal{H} , and using the norm of \mathcal{H} as a metric one can show that the distance between \hat{m} and m goes to zero almost surely under two assumptions: (i) that (S) holds, and (ii) that $d \rightarrow \infty$ and $d/n \rightarrow 0$ as $n \rightarrow \infty$. This is "best possible" in the sense that if $d/n \rightarrow c > 0$, then one cannot even get weak convergence of the distance to zero.

The second part of this project involves the estimation of the covariance, say R , of a Gaussian process whose mean is known (and may be assumed to be identically zero). In the project proposal the PI conjectured: when the Hilbert space \mathcal{H} above is infinite dimensional, the likelihood function of the covariance is unbounded, so that the maximum likelihood estimator (MLE) fails to exist.

c. The above conjecture has been proved. In particular, consider this condition:

(C) The true covariance is specified by a bivariate expansion using a fixed countable orthonormal set in \mathcal{H} .

The PI has shown that the MLE exists almost surely if the set in (C) is finite, and fails to exist (the likelihood being unbounded) almost surely if the set is infinite.

Thus a sieve estimator is necessary in general, and the "finite" case above actually leads to a natural sieve estimator \hat{R} , with dimension d as sieve parameter. The PI has established the following.

d. When (C) holds, the estimator \hat{R} , viewed as a stochastic process on TxT , has a simple distribution in terms of independent χ^2 random variables, from which it follows that, at each $(s,t) \in \text{TxT}$, $\hat{R}(s,t)$ is asymptotically unbiased as $d \rightarrow \infty$ and is weakly consistent for $R(s,t)$ if $d = O(n)$.

The coefficients in the expansion in (C) form a square-summable sequence, so we may reparametrize the model by a

subset of an ℓ^2 -space. Using the metric defined by the ℓ^2 -norm, the PI has shown the following:

e. If $d \rightarrow \infty$ and $d/n \rightarrow 0$, then we have strong consistency in ℓ^2 , under assumption (C).


f. The metric in ℓ^2 is equivalent to a natural distance between covariances given by the norm in $\mathcal{H} \otimes \mathcal{H}$, the second symmetric tensor power of \mathcal{H} .

Aside from further investigation of these sieve estimates, the next goal of the project is to weaken or remove a restriction of continuity placed on a filter for a stochastic Gaussian signal invented by M. Driscoll. Details are given in the PI's project description.

References

Geman, S., and Hwang, C.-R. (1982). Nonparametric maximum likelihood estimation by the method of sieves, Ann. Statist. 10: 401-414.

Grenander, U. (1981). Abstract Inference. Wiley, New York.



4. Written Publications:

Jay H. Beder, "A sieve estimator for the mean of a Gaussian process," Annals of Statistics (submitted).

Jay H. Beder, "A sieve estimator for the covariance of a Gaussian process," Annals of Statistics (in preparation).

5. Professional Personnel:

None besides the PI.

6. Interactions:

(a) Papers presented:

(i) "Sieve estimation for Gaussian processes," Joint Statistical Meetings, Las Vegas, Nevada, August 7, 1985.

(ii) "Sieve estimation for Gaussian processes," International Symposium on Foundations of Statistical Inference, Tel Aviv, Israel, December 15-19, 1985.

(b) Consultative and advisory functions: none.

7. New discoveries, inventions, patents: none.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

AD-A166055

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS None	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Unlimited distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 86-0069	
6a. NAME OF PERFORMING ORGANIZATION Dr. Jay H. Beder Univ. of Wisconsin-Milwaukee	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Mathematical and Information Sciences Air Force Office of Scientific Research	
6c. ADDRESS (City, State and ZIP Code) Department of Mathematical Sciences P. O. Box 413 Milwaukee, WI 53201		7b. ADDRESS (City, State and ZIP Code) Bolling Air Force Base, DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Science Corp #7	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR 84-0329	
10. SOURCE OF FUNDING NOS.			
11. TITLE (Include Security Classification) Sieves for Gaussian Processes (Unclassified)		PROGRAM ELEMENT NO. G1102F	TASK NO. A5
12. PERSONAL AUTHOR(S) Beder, Jay H.		PROJECT NO. 2304	WORK UNIT NO.
13a. TYPE OF REPORT Annual Technical Report	13b. TIME COVERED FROM 9-30-84 TO 9-29-85	14. DATE OF REPORT (Yr., Mo., Day) 11-27-85	15. PAGE COUNT 6
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
		Asymptotic unbiasedness, consistency, maximum likelihood estimation, reproducing kernel Hilbert space, sieve	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>The PI has established several important properties of a proposed sieve estimator for the mean of a Gaussian process of known covariance: The estimator is itself a Gaussian process; under a separability assumption, it is asymptotically unbiased and weakly consistent at each "time" t, and is strongly consistent (globally) in an appropriate norm.</p> <p>For a Gaussian process with zero mean and unknown covariance, the PI has shown that the likelihood for the covariance is in general unbounded almost surely. Moreover, he has developed properties of a proposed sieve estimator for the covariance analogous to those for the mean. No assumption is made about the nature of the "time" parameter t, either for the mean estimator or for the covariance estimator.</p>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Maj. Brian Woodruff		22b. TELEPHONE NUMBER (Include Area Code) 202-767-5027	22c. OFFICE SYMBOL 7111

END
FILMED

5-86

DTIC